Table 19.9 Non-Hodgkin Lymphoma

SEER^a Relative Survival (Percent) By Year of Diagnosis

All Races, Males and Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Survival Time	е																			
1-year	70.2	73.0	71.3	70.4	70.3	73.2	75.7	76.7	77.3	77.2	78.7	81.0	80.3	81.0	81.5	80.9	81.5	83.9	83.7	83.9
2-year	60.2	63.7	62.8	62.2	61.7	65.8	68.5	70.0	71.0	71.3	73.3	76.0	75.7	76.0	77.1	76.8	77.3	78.7	79.1	
3-year	54.6	58.5	58.2	57.7	57.4	62.0	64.9	66.4	66.8	67.7	70.1	73.1	73.1	73.5	74.6	74.4	74.7	75.8		
4-year	50.5	54.5	54.3	54.4	54.6	58.6	62.1	63.6	64.4	65.4	68.1	71.2	71.6	71.4	72.7	72.5	72.8			
5-year	46.8	50.7	51.0	51.6	51.9	56.0	59.4	61.0	62.0	63.7	66.0	69.3	70.0	70.0	71.3	70.3				
6-year	43.6	47.5	48.3	49.2	49.8	54.1	57.1	58.4	59.5	61.9	64.7	68.1	68.3	68.6	69.9					
7-year	40.8	45.0	45.8	46.9	47.8	52.2	55.2	56.5	57.7	60.3	62.9	66.2	67.0	66.9						
8-year	38.4	42.8	43.4	45.0	46.3	50.4	52.9	55.3	56.2	58.7	61.3	65.0	65.6							
9-year	36.5	40.8	41.4	43.4	45.2	49.3	51.5	53.6	54.7	57.3	59.9	63.9								
10-year	34.9	38.9	39.8	42.1	44.2	48.1	50.1	52.3	53.5	56.0	58.8									
11-year	33.4	37.3	38.4	40.6	43.3	46.6	48.7	51.1	52.4	54.4										
12-year	32.3	35.9	36.8	39.5	42.1	45.5	48.1	50.2	51.0											
13-year	31.3	34.7	35.7	38.3	41.3	44.7	46.8	49.2												
14-year	30.1	33.5	34.6	37.1	40.4	44.1	45.4													
15-year	28.8	32.2	33.5	36.4	39.6	43.5														
16-year	27.8	31.1	32.7	35.5	39.0															
17-year	27.3	30.3	32.0	34.8																
18-year	26.6	29.5	31.3																	
19-year	25.9	28.7	30.5																	
20-year	25.1	27.9	30.0																	

Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).